

MAT225 Section Summary: 4.3

Linearly independent sets; bases

Summary

We're accustomed to writing vectors in terms of a set of fixed vectors: for example, in two-space we write every vector in terms of vectors i and j . Each vector has a unique representation in terms of these two vectors, which is important. This set of two vectors is called a **basis** of two-space: it is enough vectors to write each vector of the space in terms of it, but not so many vectors that there are multiple representations of each vector. These are the two important properties: spanning the space, and avoiding any redundancy. That is, a basis is the smallest spanning set possible. It is also the largest set of linearly independent vectors: any more, and you'd have dependence.

linear independence: An indexed set of vectors $\{v_1, \dots, v_n\}$ in V is said to be linearly independent if the vector equation

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n = 0$$

has *only* the trivial solution ($c_1 = c_2 = \dots = c_n = 0$). The set is **linearly dependent** if it has a nontrivial solution.

Example: #4, p. 243 (independence)

$$\begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 2 \end{bmatrix}, \begin{bmatrix} -7 \\ 5 \\ 4 \end{bmatrix} \rightarrow A = \begin{bmatrix} 2 & 1 & -7 \\ -2 & -3 & 5 \\ 1 & 2 & 4 \end{bmatrix} \sim I$$

so we have independence: this is a basis.

Theorem 4: An indexed set $\{v_1, \dots, v_n\}$ of two or more vectors, $v_1 \neq 0$, is linearly dependent \iff some v_j ($j > 1$) is a linear combination of the preceding vectors $\{v_1, \dots, v_{j-1}\}$.

Example: #33, p. 245

$$p_1(t) = 1 + t^2$$

$$p_2(t) = 1 - t^2$$

$$\bar{0} \equiv p_2(t) = 0$$

1. Is $\{\bar{p}_1, \bar{p}_2\}$ a linearly independent set in \mathbb{P}_2 ?

Check for dependence: is $p_2(t) = c p_1(t)$ for some c ?

$$c p_1(t) = c(1 + t^2) = 1 - t^2 \Rightarrow$$
$$c + c t^2 = 1 - t^2 \Rightarrow$$

(setting coefficients equal) $c = 1$ and $ct^2 = -t^2$

$c = 1$ and $c = -1$

Nope!

Basis: Let H be a subspace of a vector space V . An indexed set of vectors $B = \{b_1, \dots, b_p\}$ in V is a basis for H if

1. B is a linearly independent set, and
2. the subspace spanned by B coincides with H ; that is,

$$H = \text{Span} \{b_1, \dots, b_p\}$$

Example: #4, p. 243 (basis)

Example: #34, p. 245

$$\left. \begin{array}{l} p_1(t) = 1+t \\ p_2(t) = 1-t \\ p_3(t) = 2 \end{array} \right\} \text{dependent: } \bar{p}_1 + \bar{p}_2 = \bar{p}_3$$

$\{\bar{p}_1, \bar{p}_2\}$ is a basis for $\text{Span}\{\bar{p}_1, \bar{p}_2, \bar{p}_3\}$ (checked linear independence). Other choices: $\{\bar{p}_1, \bar{p}_3\}$, $\{\bar{p}_2, \bar{p}_3\}$, $\{6\bar{p}_1, 3\bar{p}_2\}$, etc.

Example: The columns of the $n \times n$ identity matrix I_n form a basis, called the **standard basis** for \mathbb{R}^n :

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

$$\hat{i} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad \hat{j} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad \hat{k} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

In three-space these are simply the vectors \mathbf{i} , \mathbf{j} , and \mathbf{k} .

Theorem 5 (the spanning set theorem): Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$ be a set in V , and let $H = \text{Span} \{\mathbf{v}_1, \dots, \mathbf{v}_p\}$.

1. If one of the vectors in S – say \mathbf{v}_k – is a linear combination of the remaining vectors in S , then the set formed from S by removing \mathbf{v}_k still spans H .
2. If $H \neq \{0\}$, some subset of S is a basis for H .

Theorem 6: the pivot columns of a matrix A form a basis for $\text{Col } A$.

Turns out that elementary row operations on a matrix do not affect the linear dependence relations among the columns of the matrix. Hence, the reduced matrix has the same independent columns as the original matrix. Make sure to choose the columns of the matrix A , however, rather than the reduced matrix....

Example: #36, p. 245