

Section 11.1: Sequences

Review

Questions

- Evaluate $\int_1^{\infty} \frac{1}{x^\pi} dx$ and $\int_1^{\infty} \frac{1}{x^{0.1}} dx$.
- For which exponents p does $\int_1^{\infty} \frac{1}{x^p} dx$ converge?

Questions

- Evaluate $\int_0^1 \frac{1}{x^\pi} dx$ and $\int_0^1 \frac{1}{x^{0.1}} dx$.
 - For which exponents p does $\int_0^1 \frac{1}{x^p} dx$ converge?
-

Approximations

Questions

Use the trapezoid rule to approximate $\int_0^1 \cos(\sqrt{x}) dx$

- using $n = 1$
- using $n = 2$
- using $n = 3$
- using $n = 4$
- What do we expect to happen as we use larger and larger values of n in the trapezoid rule?

Questions

A standard way to approximate the square root of a value is as follows. This method was known to the Babylonians, a few thousand years ago.... To approximate \sqrt{a}

1. Make a rough estimate for the value of \sqrt{a} . Call it the first approximation x_1 .
2. The second approximation is $x_2 = 0.5 \left(x_1 + \frac{a}{x_1} \right)$.
3. The third approximation is $x_3 = 0.5 \left(x_2 + \frac{a}{x_2} \right)$.
4. In general, the n^{th} approximation is $x_n = 0.5 \left(x_{n-1} + \frac{a}{x_{n-1}} \right)$.

- Use this scheme to find the fourth approximation to $\sqrt{2}$ using $x_1 = 2$.
- Do the approximations improve as n gets larger?

This method is an example of “Newton’s method” (section 3.8), which is a very general method for finding roots (zeros) of functions.

Sequences

Definition

A *sequence* is an infinite list of numbers written in a definite order

$$a_1, a_2, a_3, \dots, a_k, \dots$$

Examples

- 1, 2, 3, 4, 5, ...
- $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$
- -1, 2, -3, 4, -5, 6, -7, 8, ...

Forms

Besides writing a sequence as a list we will also write some sequences

- as formulas: for example $\left\{\frac{n}{2^n}\right\}_{n=1}^{\infty}$
- recursively: for example $f_1 = 1, f_2 = 1, f_n = f_{n-1} + f_{n-2}$

Limits

Frequently, but not always, sequences $\{a_n\}$ are lists of approximations that converge to some desired exact values.

$$\lim_{n \rightarrow \infty} a_n$$

Questions

Do any of the above sequences converge? If so, to what values?

- $x_1 = 2, x_n = 0.5 \left(x_{n-1} + \frac{a}{x_{n-1}} \right)$
- $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots$
- -1, 2, -3, 4, -5, 6, -7, 8, ...
- $1, -\frac{1}{4}, \frac{1}{9}, -\frac{1}{16}, \frac{1}{25}, -\frac{1}{36}, \dots$
- $\left\{\frac{n}{2^n}\right\}_{n=1}^{\infty}$

- $f_1 = 1, f_2 = 1, f_n = f_{n-1} + f_{n-2}$

Techniques

Technique 1

If $a_n = f(n)$ where f is a real-valued function, then $\lim_{n \rightarrow \infty} a_n = \lim_{x \rightarrow \infty} f(x)$.

Questions

- Find $\lim_{n \rightarrow \infty} \frac{n}{n^2+1}$ by evaluation the limit $\lim_{x \rightarrow \infty} \frac{x}{x^2+1}$.
- For which of the above sequences can you use this technique?

Technique 2

If $\lim_{n \rightarrow \infty} |a_n| = 0$, then $\lim_{n \rightarrow \infty} a_n = 0$.

Question

- Does this technique apply to any of the above limits?
- For which values of r does the sequence $\{r^n\}$ converge?

Monotonic sequences

Questions

- What should we mean by an *increasing* sequence?
- What is an example of an increasing sequence?
- What should we mean by a *decreasing* sequence?
- What is an example of a decreasing sequence?
- What is an example of a sequence that is neither increasing nor decreasing?

Definition

A sequence that is either increasing or decreasing is said to be monotonic.

Bounded sequences

- What should we mean by a sequence that is *bounded above*?
- What is an example of a sequence that is bounded above?
- What should we mean by a sequence that is *bounded below*?

- What is an example of a sequence that is bounded below?

Technique 3

A bounded, monotonic sequence converges.

You might think this way: *the sequence has to go somewhere -- it's constantly increasing (say), and yet it can't go past a certain point -- so it has to go somewhere between where it is at any moment and that upper bound.* Furthermore, this is true even as long as the sequence is **eventually** monotonic.

You might think this way: *no finite chunk of a sequence at the beginning has any impact on convergence. Convergence is a property of the long tail, as the index sails off to infinity.*

Questions

Consider the sequence $\left\{\frac{n}{2^n}\right\}_{n=1}^{\infty}$.

- Is this sequence monotonic?
- Is this sequence bounded?